

# Validation of a food-frequency questionnaire for measurement of nutrient intake in a dietary intervention study

Gina Segovia-Siapco<sup>1,2,\*</sup>, Pramil Singh<sup>3</sup>, Karen Jaceldo-Siegl<sup>1,4</sup> and Joan Sabaté<sup>1,3,4</sup>

<sup>1</sup>Department of Nutrition, School of Public Health, Loma Linda University, Loma Linda, CA, USA: <sup>2</sup>Department of Public Health, School of Graduate Studies, Adventist International Institute of Advanced Studies, PO Box 038, Silang 4118, Cavite, Philippines: <sup>3</sup>Department of Epidemiology and Biostatistics, School of Public Health, Loma Linda University, Loma Linda, CA, USA: <sup>4</sup>The Adventist Health Study II, School of Public Health, Loma Linda University, Loma Linda, CA, USA

Submitted 23 December 2005; Accepted 23 May 2006

## Abstract

**Objective:** To validate a 171-item semi-quantitative food-frequency questionnaire (FFQ) for measurement of nutrient intake in an intervention trial based on walnut supplementation.

**Design and setting:** Free-living adults from Southern California were randomly assigned to either an intervention (walnut-supplemented) or a control diet. The prescribed 6-month intervention was  $\geq 28$  g of walnuts per day for the walnut-supplemented group and  $\leq 2$  g of walnuts per day for the control group. Participants provided at least six 24-hour dietary recalls and completed a self-administered FFQ.

**Subjects:** Eighty-seven adults aged 30–72 years (48 females, 39 males).

**Results:** Our findings from validation (by correlation with six diet recall measures) of the measurement of 32 nutrients by the FFQ are as follows. We found significant positive correlations (corrected for measurement error) between the FFQ and diet recalls for total energy ( $r = 0.34$ ), total carbohydrate ( $r = 0.42$ ), vegetable protein ( $r = 0.43$ ), total fat ( $r = 0.51$ ), polyunsaturated fat ( $r = 0.77$ ), total fibre ( $r = 0.60$ ), linoleic acid ( $r = 0.78$ ) and  $\alpha$ -linolenic acid ( $r = 0.79$ ) – the last nutrient being an excellent nutrient biomarker of the intervention (walnut supplementation). Significant positive correlations were also found for vitamin C ( $r = 0.96$ ) and certain minerals ( $r = 0.46$ – $0.80$  for calcium, phosphorus, magnesium, iron and potassium). Uncorrected correlations were also high ( $r > 0.40$ ) for retinol,  $\beta$ -carotene, folate and alcohol. Both diet recalls and FFQ showed a similar significant difference in  $\alpha$ -linolenic acid content between the walnut-supplemented and control diets.

**Conclusions:** The FFQ demonstrated good relative validity in the estimation intake of some of the major nutrients in a dietary intervention trial and was a particularly valid estimate of an important nutrient biomarker of walnut supplementation.

## Keywords

Validity  
Food-frequency questionnaire  
24-hour dietary recalls  
Nutrient intake  
Intervention  
Free-living

Dietary modification is considered an important component in the prevention, management and treatment of chronic conditions such as hypertension<sup>1</sup>, hypercholesterolaemia<sup>2</sup>, diabetes<sup>3</sup> and obesity<sup>4</sup>. Recently, there have been a number of large-scale clinical trials that continue to test the effect of dietary modification and/or supplementation on primary and secondary prevention of a wide range of chronic and infectious diseases<sup>5–8</sup>.

In large-scale community-based intervention trials, the more precise methods of dietary assessment such as interviewer-based dietary recalls, food records or lengthy diet histories can be prohibitively expensive and carry a respondent burden that contributes to a high drop-out rate – particularly in underserved communities<sup>9</sup>. A practical and cost-effective alternative to diet recalls and diet histories is to assess dietary intake with a self-administered

food-frequency questionnaire (FFQ). Numerous validation studies have shown that in a prospective cohort study, the FFQ can be a valid estimate of the food and nutrients measured by dietary recalls or diet histories<sup>10</sup>. To date, only a few studies have validated the use of the FFQ to assess baseline diet and dietary change (due to prescribed intervention) in an intervention trial<sup>11–15</sup>.

In the present study, we have conducted a validation analysis of data from a 6-month intervention trial that investigated the health effect of daily incorporation of walnuts into the habitual diet of free-living individuals. Dietary data were collected at baseline and during the follow-up by multiple 24-hour dietary recalls and by a self-administered 171-item semi-quantitative FFQ. The purposes of the present validation study were (1) to examine whether the FFQ provided a valid measure of nutrient

intake among subjects enrolled in a clinical trial, and (2) to examine whether the FFQ can measure the effect of dietary intervention (walnut supplementation) on intake of specific nutrients. Findings from this study will contribute to the design of an FFQ that can be used in large-scale community intervention trials of diet and health outcomes.

## Materials and methods

### Study population

As previously described<sup>16</sup>, the study population was recruited from 94 volunteers who responded to local advertisements from Loma Linda University about a feeding trial. Three of the 94 participants dropped out due to compliance difficulty, one did not fill out the FFQ and three completed fewer than six dietary recalls. Thus, the analytical population for this validation study consisted of 87 subjects who completed at least six dietary recalls and the self-administered FFQ. Among the recalls collected from the 87 subjects, five recalls were excluded due to their extreme total energy values. We defined 'extreme' as being outside the biologically plausible range of 800–4500 kcal among men and outside the biologically plausible range of 500–3500 kcal among women.

### Study design

At baseline, subjects were randomly assigned to either the control (habitual or usual diet) or the walnut-supplemented group. The control group was instructed to eat their usual diet and refrain from eating walnuts ( $\leq 2$  g of walnuts per day) and substantial amount of other nuts, including nut butters, during the study. The intervention group was provided with 25–56 g of pre-packaged raw walnuts to be eaten daily together with their usual diet in any way they preferred. This amount of walnuts accounted for  $\sim 12\%$  of the subject's total daily energy intake. Since the intervention was among free-living adults, we did not give any other dietary advice to the participants. During the 6-month follow-up, we collected, through telephone interviews, at least six 24-hour dietary recalls (mean = 6.9 recalls) at intervals of 2–5 weeks. The protocol for these recalls is given below. At the end of 6 months, the subjects completed a self-administered FFQ which assessed intake during the past 6 months.

### FFQ

We modified a previously validated FFQ that was developed at our institution<sup>17,18</sup>. The original questionnaire was designed to allow a measure of vegetarian diet patterns (i.e. meat analogues). As described previously<sup>17,18</sup> for the intervention trial in a general population sample, we removed many of the items on specific meat analogues and added additional items on meat intake. The questionnaire was then pilot-tested for clarity, interpretation and improvement of format in a

substudy of 10 individuals, who had similar demographic characteristics to the study group but were not participants in the study.

The FFQ is semi-quantitative and consists of 171 hard-coded food and 15 write-in items. The respondent is asked to report average consumption for the past 6 months. Food items are grouped under the following categories: breads, grains and starches; vegetables; legumes and nuts; eggs, dairy products, oils; fish and meats; fruits; beverages; sweets and baked goods; and condiments, dressings, miscellaneous. Portion sizes are based on average serving sizes using familiar measuring devices, e.g. cup, teaspoon, can and others. The frequency section consists of eight categories: never or rarely, 1–3 per month, 1 per week, 2–4 per week, 5–6 per week, 1 per day, 2–3 per day and 4+ per day. Except for four write-in items (cold breakfast cereals), lack of substantial respondent information precluded inclusion of 11 write-in items in the validation of this questionnaire.

### Reference method: 24-hour dietary recalls

Research nutritionists collected 24-hour dietary recalls during telephone interviews with the participants using a protocol that has been previously described<sup>16</sup>. The research nutritionists trained in the use of Nutrition Data Systems (NDS-R) for Research software package<sup>19</sup>. The NDS-R system enables collection of data on 18 000 foods, 8000 brand name products and many ethnic foods. For 24-hour recalls, the NDS software has built-in quality control with a multiple-pass approach interview methodology<sup>19</sup>. Default food description assignments on the software standardise data collection and therefore minimise inter-interviewer variation. A two-dimensional food portion visual was provided to help participants estimate food portion sizes.

The telephone interviews were unannounced, unscheduled, and included a standard script with instructions that were read by each research nutritionist before the NDS-R protocol was implemented. On average, recall interviews lasted 30 min. Each participant provided at least six dietary recalls (mean  $\pm$  standard deviation (SD) for recalls =  $6.9 \pm 0.4$ ). Day-to-day variation was accounted for by obtaining recalls on either (1) all days of the week or (2) at least one weekend day and five weekdays.

### Data analyses

#### Validation

Collection and coding of dietary intakes on both the FFQ and dietary recalls were performed using NDS-R software. To determine average frequency of intake per day on the FFQ, we converted the frequency value of average consumption over the past 6 months into frequency of intake per day (e.g. frequency of 1–3 per month =  $2/30 = 0.067$ , and so on). The factors were 0 for never or rarely, 0.067 for 1–3 per month, 0.143 for 1 per

week, 0.429 for 2–4 per week, 0.786 for 5–6 per week, 1.0 for 1 per day, 2.5 for 2–3 per day, and 5 for 4+ per day. Nutrient amounts were relative to the fixed portion/serving sizes in the semi-quantitative FFQ. For multiple-food items, the nutrient compositions of the more commonly eaten varieties of the foods in the item were averaged, e.g. the nutrient composition computed for a multiple food-item that consists of roll, English muffin, bagel or soft pretzel was based on the average of the nutrient compositions of a medium size (white flour) roll, medium size English muffin, medium size (white flour) bagel and medium size soft pretzel. The average of raw and cooked nutrient components of vegetables known to be eaten either raw or cooked was used for vegetables such as carrots, broccoli and others. For single food items where several types exist, the NDS-R software computes nutrient composition for the most commonly eaten type – a determination made using nationally representative market research data<sup>19</sup>. Nutrient intake per day was then computed using the sum-product method, i.e.

$$A = \sum_{i=1}^n F_i a_i,$$

where  $A$  = total intake of nutrient  $A$  per day;  $F_i$  = frequency of food  $i$  intake per day; and  $a_i$  = amount of nutrient  $A$  in food  $i$ .

To determine the relative validity of the FFQ's nutrient intake estimates at the group level, we compared its means and standard deviations with those of the reference method using paired samples  $t$ -test. For relative validity at the individual level, we performed correlation analyses. Since our intent was to determine the amount of variation explained by the test method on the reference method (i.e.  $R^2$ ), the use of parametric (Pearson's) correlation analyses does not assume any particular underlying distribution of the variables<sup>20</sup>. To correct for within-person variation in multiple dietary intake measurements with the reference method (dietary recalls), the following formula was used<sup>10</sup>:

$$r_c = r_0 \sqrt{1 + \frac{S_w^2/S_b^2}{n}},$$

where  $r_c$  = corrected/de-attenuated correlation coefficient;  $r_0$  = uncorrected/attenuated correlation between FFQ and multiple 24-hour recalls;  $S_w^2$  = within-person variance of the multiple 24-hour recalls;  $S_b^2$  = between-person estimate of variance in the reference method (24-hour recalls); and  $n$  = number of repeated measures of the dietary recalls.

De-attenuation of the correlation coefficients using the method described above creates conditions in which we could not assume normally distributed errors for  $r_c$ . Thus, instead of the traditional asymptotic methods to determine confidence intervals about  $r_c$ , we computed 'distribution-free', non-parametric 95% confidence inter-

vals using the BCa bootstrap re-sampling method<sup>21</sup>, with each confidence interval determined from the distribution of  $r_c$ s from 2000 samples.

Nutrient variables were also adjusted for energy since amounts of walnuts allotted to the participants differed according to their energy intake. Nutrient intake distributions were normalised by logarithmic and square root transformations before using the residual method for energy adjustment. Energy-adjusted correlations were obtained using Pearson's correlations.

#### Intervention measure

We were interested in finding out if the FFQ would be able to assess the prescribed intervention measure in the two diet groups. The test food, walnut, is high in  $\alpha$ -linolenic acid (ALA) (9.1 g/100 g walnuts) and linoleic acid (LA) (38.1 g/100 g walnuts). Walnut studies have shown increases in plasma levels of LA and ALA as a result of incorporation of walnuts into the diet<sup>22,23</sup>. Although the amount of LA in walnuts is much higher compared with that of ALA, LA is ubiquitous in the American diet and using it as a marker of intervention measure in this study may be confounded by other food sources of LA.

ALA is a more unique component of walnuts than LA. It is important to note that ALA is also found in items such as flaxseed, flaxseed oil, canola oil and soybean oil. We did not expect that walnut supplementation would substantially change intake of these items.

To assess further whether ALA would be the most appropriate marker of the intervention measure, we determined the percentage of energy accounted for by both LA and ALA, using 1.1% as the cut-off for ALA (i.e. the percentage of energy accounted for by ALA in 28 g of walnuts if the total energy requirement is 2000 kcal day<sup>-1</sup>) and 4.8% as the cut-off for LA (i.e. the percentage of energy accounted for by LA in 28 g of walnuts eaten if the total energy requirement is 2000 kcal/day<sup>-1</sup>). We then compared the calculated percentage of energy values for each diet group with this criterion. Reference method (dietary recall) values of mean percentage energy from ALA intake were 2.2% (SD = 0.4) for the walnut diet group and 0.6% (SD = 0.2) for the control diet group. For LA, however, reference method values of mean percentage energy from LA intake were 11.3% (SD = 1.9) for the walnut study group and 5.6% (SD = 1.6) for the control diet group. Determining a cut-off value for LA that will delineate intervention from non-intervention was more challenging than for ALA because LA is not unique to walnuts. Our use of ALA as the nutrient marker for the intervention is, therefore, justified.

We assumed that proportional increases in ALA intake would be a consequence of consuming 28–56 g of walnuts. As already shown earlier, for an individual whose daily energy requirement is 2000 kcal, 28 g of walnuts will result in an ALA intake that accounts for ~1.1% of that individual's daily energy requirement.

Utilising this logic, we defined the intervention measure as ALA intake  $\geq 1.1\%$  energy for the walnut diet group and ALA intake  $< 1.1\%$  of energy for the control diet group.

Means of absolute estimates ( $\pm$ SD) of ALA by the FFQ and the 24-hour dietary recalls were computed for both diet groups. We used cross-classification to determine agreement between the two methods in assessing adherence to the prescribed intervention measure described above.


Graphical approaches were also used to determine agreement between the test and reference methods in the estimation of ALA – the nutrient most indicative of the intervention effect. Specifically, we plotted the FFQ and recall values of ALA in the control and walnut group (Fig. 1). We also constructed a Bland–Altman plot to assess agreement between the methods by plotting the difference between FFQ and recalls vs. the recall values.

## Results

Demographic characteristics of the study population of 87 subjects are given in Table 1 and are further stratified by intervention group. The study population was 74% Caucasian and was aged 30–72 years (mean = 54.7). We found no important demographic differences between the walnut-supplemented and control groups.

### Validation of FFQ nutrient measures in all subjects

In Table 2 we have reported the mean values for 32 nutrients computed from the FFQ and the corresponding nutrients computed from the mean of the diet recalls. Mean nutrient intake on the FFQ was higher than dietary recalls for all nutrients, and these differences were statistically significant for all nutrients except for animal protein, ALA, cholesterol and alcohol. Total energy intake was overestimated by 160 kcal (8.4%), and for the



**Fig. 1** Plot of  $\alpha$ -linolenic acid (ALA) intake estimate values from the food-frequency questionnaire (FFQ) against dietary recall estimates of ALA intake

**Table 1** Demographic characteristics of study subjects

Characteristic	By diet groups		
	All subjects ( <i>n</i> = 87)	Walnut ( <i>n</i> = 48)	Control ( <i>n</i> = 39)
Females (%)	55	58	51
Recalls (no.)	6.9 (0.4)	6.9 (0.4)	6.8 (0.5)
Age (years)	54.7 (10.3)	55.3 (9.9)	54.0 (10.7)
Height (cm)	169.0 (10.4)	167.6 (8.9)	170.7 (11.9)
Weight (kg)	75.7 (14.1)	73.2 (13.2)	78.7 (14.8)
BMI (kg m <sup>-2</sup> )	26.4 (3.4)	26.0 (3.5)	26.9 (3.2)

BMI – body mass index.

All values are reported as mean (standard deviation).

Mean height and mean weight are significantly different between males and females.

remaining nutrients the overestimation ranged from 1.3% (retinol) to 87.5% (docosahexaenoic acid).

Table 3 provides the de-attenuated (corrected for measurement error in the 24-hour diet recalls) correlations for 19 out of the 32 nutrients measured by the FFQ and the mean of at least six 24-hour dietary recalls. Statistically

**Table 2** Nutrient intake estimates of the food-frequency questionnaire (FFQ) and 24-hour dietary recalls (DR) for all participants (*n* = 87)

Nutrient	Mean $\pm$ SD	
	FFQ	DR
Energy (kcal)	2073 $\pm$ 711*	1913 $\pm$ 609
Total CHO (g)	262.0 $\pm$ 87.8*	245.2 $\pm$ 86.8
Total protein (g)	75.7 $\pm$ 37.3*	72.3 $\pm$ 26.0
Animal protein (g)	45.2 $\pm$ 32.0	43.5 $\pm$ 21.0
Vegetable protein (g)	30.0 $\pm$ 12.0*	28.2 $\pm$ 11.3
Total fat (g)	85.2 $\pm$ 37.8*	74.2 $\pm$ 26.0
Total SFA (g)	23.6 $\pm$ 12.8*	21.7 $\pm$ 8.9
Total MUFA (g)	30.4 $\pm$ 12.9*	25.1 $\pm$ 9.7
Oleic acid (g)	28.6 $\pm$ 12.0*	23.6 $\pm$ 9.2
Total PUFA (g)	25.0 $\pm$ 13.0*	21.9 $\pm$ 9.7
LA (g)	21.3 $\pm$ 10.9*	18.5 $\pm$ 8.1
ALA (g)	3.3 $\pm$ 2.1	3.1 $\pm$ 1.8
Arachidonic acid (mg)	0.13 $\pm$ 0.13*	0.10 $\pm$ 0.06
EPA (mg)	0.06 $\pm$ 0.06*	0.04 $\pm$ 0.05
DHA (mg)	0.15 $\pm$ 0.15*	0.08 $\pm$ 0.10
Cholesterol (mg)	212.5 $\pm$ 158.5	205.2 $\pm$ 101.4
Alcohol (g)	3.3 $\pm$ 5.9	3.0 $\pm$ 5.8
Total dietary fibre (g)	25.4 $\pm$ 9.4*	20.6 $\pm$ 6.6
$\beta$ -Carotene ( $\mu$ g)	5090 $\pm$ 3978*	3943 $\pm$ 3539
Retinol ( $\mu$ g)	472 $\pm$ 300*	466 $\pm$ 446
Total vitamin E (mg)	14.0 $\pm$ 5.5*	9.2 $\pm$ 3.7
$\alpha$ -Tocopherol (mg)	8.9 $\pm$ 3.5*	7.0 $\pm$ 2.7
$\beta$ -Tocopherol (mg)	0.4 $\pm$ 0.2*	0.3 $\pm$ 0.1
$\gamma$ -Tocopherol (mg)	19.1 $\pm$ 9.2*	16.2 $\pm$ 7.0
Vitamin C (mg)	152.0 $\pm$ 94.0*	100.4 $\pm$ 48.8
Folate ( $\mu$ g)	454 $\pm$ 170*	401 $\pm$ 156
Calcium (mg)	942 $\pm$ 565*	801 $\pm$ 429
Phosphorus (mg)	1375 $\pm$ 600*	1229 $\pm$ 471
Magnesium (mg)	356.6 $\pm$ 120.2*	322.6 $\pm$ 102.3
Iron (mg)	16.5 $\pm$ 7.3*	15.3 $\pm$ 6.4
Zinc (mg)	10.6 $\pm$ 4.8*	10.3 $\pm$ 5.2
Potassium (mg)	3359 $\pm$ 1269*	2818 $\pm$ 896

SD – standard deviation; CHO – carbohydrate; SFA – saturated fatty acid; MUFA – monounsaturated fatty acid; PUFA – polyunsaturated fatty acid; LA – linoleic acid; ALA –  $\alpha$ -linolenic acid; EPA – eicosapentaenoic acid; DHA – docosahexaenoic acid.

\*Significantly different from DR values at  $P < 0.0001$ .

**Table 3** De-attenuated (corrected for measurement error in the 24-hour diet recall standard) correlation coefficients between the food-frequency questionnaire and 24-hour dietary recall (at least six recalls per subject) measures of 19 nutrients are presented with non-parametric confidence limits (CI)

Nutrient	<i>r</i>	95% CI
Energy (kcal)	0.39	0.04, 0.63
Total CHO (g)	0.42	0.13, 0.64
Total protein (g)	0.33	-0.36, 0.67
Animal protein (g)	0.36	*
Vegetable protein (g)	0.43	0.09, 0.74
Total fat (g)	0.51	0.13, 0.87
Total SFA (g)	0.59	-0.11, 1.00
Total MUFA (g)	0.39	-0.17, 1.00
Oleic acid (g)	0.43	-0.19, 1.00
Total PUFA (g)	0.77	0.54, 0.89
LA (g)	0.78	0.51, 0.92
ALA (g)	0.79	0.55, 0.92
Total dietary fibre (g)	0.60	0.21, 0.89
Vitamin C (mg)	0.96	0.40, 1.00
Calcium (mg)	0.70	0.46, 0.81
Phosphorus (mg)	0.46	0.11, 0.61
Magnesium (mg)	0.55	0.25, 0.73
Iron (mg)	0.60	0.18, 0.94
Potassium (mg)	0.48	0.21, 0.70

CHO – carbohydrate; SFA – saturated fatty acid; MUFA – monounsaturated fatty acid; PUFA – polyunsaturated fatty acid; LA – linoleic acid; ALA –  $\alpha$ -linolenic acid.

\* The confidence interval could not be computed due to high between-person variance.

significant de-attenuated correlations were found for total energy ( $r = 0.39$ ), total carbohydrate ( $r = 0.42$ ), vegetable protein ( $r = 0.43$ ), total fat ( $r = 0.51$ ), polyunsaturated fat ( $r = 0.77$ ), LA ( $r = 0.78$ ), ALA ( $r = 0.79$ ), and total fibre ( $r = 0.60$ ). The de-attenuated correlation for saturated fat was high ( $r = 0.59$ ), but did not attain significance. For the vitamins and minerals, a very high de-attenuated correlation was found for vitamin C ( $r = 0.96$ ), and significant correlations in the range of 0.46–0.80 were found for calcium, phosphorus, magnesium, iron and potassium.

Table 4 provides uncorrected correlation coefficients for the remaining 13 nutrients that could not be corrected for measurement error due to very high between-person variance. Among these 13 nutrients, strong correlations ( $>0.40$ ) were found for retinol,  $\beta$ -carotene, folate and alcohol. Further adjustment for total energy intake tended to strengthen the positive correlations for seven out of the 13 nutrients and, after energy adjustment, all attained statistical significance.

### Intervention measure

As shown in Table 3, ALA, a good nutrient marker for walnut supplementation, was measured well by the FFQ, with a correlation of 0.79 with the diet recalls. In the trial, we had used the following ALA-based criterion to distinguish between the two intervention groups: ALA was  $\geq 1.1\%$  of total energy for the walnut-supplemented group and  $< 1.1\%$  of total energy for the controls. In Table 5 we present the percentage total energy from ALA for both

**Table 4** Correlation coefficients between the food-frequency questionnaire and the 24-hour dietary recall (at least six recalls per subject) measures of 13 nutrients are presented with and without adjustment for total energy intake

Nutrient	<i>r</i>	Energy-adjusted
		<i>r</i>
Arachidonic acid (mg)	0.14	0.36**
EPA (mg)	0.20	0.26*
DHA (mg)	0.15	0.25*
Cholesterol (mg)	0.22*	0.54***
Alcohol (g)	0.84***	0.80***
$\beta$ -Carotene ( $\mu$ g)	0.69***	0.53***
Retinol ( $\mu$ g)	0.46***	0.42***
Total vitamin E (mg)	0.08	0.22*
$\alpha$ -Tocopherol (mg)	0.09	0.23*
$\beta$ -Tocopherol (mg)	0.31**	0.24*
$\gamma$ -Tocopherol (mg)	0.36**	0.50***
Folate ( $\mu$ g)	0.48***	0.39**
Zinc (mg)	0.33**	0.31**

EPA – eicosapentaenoic acid; DHA – docosahexaenoic acid.

Correction for measurement error in the 24-hour recalls was not possible for these nutrients due to high between-person variation.

\* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.0001$ .


groups that was measured by the FFQ and diet recalls. Both the FFQ and the diet recalls found a significantly higher percentage of total energy from ALA in the walnut-supplemented group ( $P < 0.05$  for walnut-supplemented vs. control on both FFQ and diet recall). We also examined the sensitivity of the FFQ in classifying subjects into the intervention groups based on being  $\geq 1.1\%$  and  $< 1.1\%$  total energy from ALA in their diet recalls. We found that in the controls, the sensitivity of the FFQ in detecting  $< 1.1\%$  total energy from ALA was 80% and in the walnut-supplemented group the sensitivity of the FFQ in detecting  $\geq 1.1\%$  total energy from ALA was 98%.

Figures 1 and 2 show that the FFQ estimates closely paralleled the recall values in their indication that ALA intake in the walnut-supplemented group was greater than that in the control group. The Bland–Altman plot (Fig. 2) indicates that there was good agreement between the two methods as evidenced by the symmetry of the data around the zero value (indicating a zero difference between the FFQ and recall measures of ALA). Taken together, these findings indicate that the FFQ was a good measure of the nutrient effect (i.e. increased ALA intake) of the walnut-supplemented intervention.

**Table 5** Percentage energy from  $\alpha$ -linolenic acid (ALA) for the walnut-supplemented diet and control that were measured by the food-frequency questionnaire (FFQ) and 24-hour dietary recalls (DR)

Group	% total energy from ALA intake (mean $\pm$ SD)	
	FFQ	DR
Walnut diet ( $n = 48$ )	1.9 $\pm$ 0.5	2.2 $\pm$ 0.4
Control diet ( $n = 39$ )	0.8 $\pm$ 0.3	0.6 $\pm$ 0.2

SD – standard deviation.



**Fig. 2** Difference between the  $\alpha$ -linolenic acid (ALA) intakes estimated from the food-frequency questionnaire (FFQ) and the reference method (24-hour dietary recall) plotted against the mean of the two methods. The solid line represents the mean while the dotted line represents the limits of agreement (95% confidence interval)

## Discussion

We sought to validate the measurement of 32 nutrients by a 171-item FFQ that was administered during a dietary intervention trial among free-living individuals. After correction for measurement error in the standard (multiple diet recalls during the follow-up), we found significant positive correlations between the FFQ and diet recalls for total energy ( $r = 0.39$ ), total carbohydrate ( $r = 0.42$ ), vegetable protein ( $r = 0.43$ ), total fat ( $r = 0.51$ ), polyunsaturated fat ( $r = 0.77$ ), total fibre ( $r = 0.60$ ), LA ( $r = 0.78$ ) and ALA ( $r = 0.79$ ) – the last nutrient being an excellent biomarker of the intervention with walnut supplementation. Among the nutrient correlations that could not be corrected for measurement error due to high between-person variance, we also found good validity (uncorrected correlations  $> 0.40$ ) for retinol,  $\beta$ -carotene, folate and alcohol. Energy adjustment tended to increase the magnitude and statistical significance of all uncorrected correlations.

Absolute intake estimates on the FFQ were significantly higher (up to 88%) for most of the nutrients compared with the reference method estimates. The length of the questionnaire, wherein a number of food items might not have been consumed or reported to have been consumed during any of the dietary recalls, could have contributed to this bias. While results vary, FFQs usually have higher estimates for most nutrients than the reference method<sup>24,25</sup> particularly if the FFQ exceeds 100 items<sup>26–28</sup>. Overestimation in dietary assessment questionnaires may be due to measurement errors introduced by differences in conceptualisation of portion sizes, misinterpretation of specific items<sup>10</sup>, and frequency and serving size differences between the test and reference methods. This is especially the case when there are multiple foods in an

item<sup>29</sup>. Moreover, averaging amounts of intake over a long period of time, as with an FFQ, is subject to estimation misjudgement that may be different from estimation errors in reporting amounts of intake over the past few hours, as in dietary recalls.

The lower, albeit significant, correlation of 0.39 for total energy is noteworthy. This may be consistent with other studies indicating that subjects tend to under-report their total energy intake on diet recalls administered during intervention trials<sup>30</sup>. In our study, the FFQ values tended to be  $\sim 160$  kcal more than the diet recalls. Further investigation of under-reporting in the recalls would require validation of the standard to values from direct measurement such as the doubly labelled water technique. We examined under-reporting of total energy by comparing total energy values with change in body weight during the first 6 months of the trial<sup>31</sup> and found possible under-reporting of values. Using Goldberg's method, 38% of the participants were classified as under-reporters, and 40% of the participants who reported an energy intake lower than their estimated energy expenditure actually gained  $> 0.50$  kg body weight. We found that exclusion of under-reporters from the analysis did not substantially change the magnitude of the correlations.

Correcting for random errors associated with within-person variation almost always increases correlation values<sup>32</sup>, and our results for de-attenuated correlations follow this same trend. However, high within-to-between person variances resulted in very high correction factors for 13 of the 32 nutrients and therefore in Table 3 we reported only uncorrected correlations. Further studies with a larger sample size are needed to investigate corrected correlations for these 13 nutrients. Another noteworthy factor is the possibility of errors in the estimation of the within-person and between-person variances<sup>10</sup>.

The ability of the FFQ to detect the intervention measure (walnut supplementation) among the diet groups was clearly shown by the high correlations with ALA in Tables 3 and 5. Walnuts are unique as a nut because of their ALA (18:3n-3) content. Both diet groups were instructed at the beginning of the study to eat or to not eat walnuts according to their diet assignment. Thus, the possibility that a training effect and compliance bias may have contributed to the relatively high agreement between FFQ and dietary recall regarding ALA intake for both diet groups cannot be discounted. It is important to note that our study examined the correlation between diet recalls and FFQ values during the intervention trial. Further studies are needed to determine whether the change in nutrient intake between baseline and follow-up is detected by the FFQ.

Our findings indicate that the FFQ did demonstrate excellent relative validity (compared with recalls) in the estimation of specific fatty acids (i.e. ALA) associated with a single food intervention (walnut), but had less relative validity in the estimation of macronutrients (i.e. total

protein and saturated fat). Further work is needed to determine whether the FFQ can estimate nutrient intake in trials where the intervention diet is complex (i.e. a diet pattern) and the treatment effect represents a change in macronutrient intake (i.e. change in protein).

### Acknowledgements

We would like to acknowledge the research nutritionists who helped collect the 24-hour dietary recalls and the California Walnut Commission for the funding of this research.

### References

- Moore TJ, Conlin PR, Ard J, Svetkey LP. DASH (Dietary Approaches to Stop Hypertension) diet is effective treatment for stage 1 isolated systolic hypertension. *Hypertension* 2001; **38**: 155–8.
- Ammerman AS, Keyserling TC, Atwood JR, Hosking JD, Zayed H, Krasny C. A randomized controlled trial of a public health nurse directed treatment program for rural patients with high blood cholesterol. *Preventive Medicine* 2003; **36**: 340–51.
- Steyn NP, Mann J, Bennett PH, Temple N, Zimmet P, Tuomilehto J, *et al.* Diet, nutrition and the prevention of type 2 diabetes. *Public Health Nutrition* 2004; **7**(1A): 147–65.
- Nicklas BJ, Dennis KE, Berman DM, Sorkin J, Ryan AS, Goldberg AP. Lifestyle intervention of hypocaloric dieting and walking reduces abdominal obesity and improves coronary heart disease risk factors in obese, postmenopausal, African-American and Caucasian women. *Journal of Gerontology Series A, Biological Sciences and Medical Sciences* 2003; **58**: 181–9.
- Bazzano LA, Serdula M, Liu S. Prevention of type 2 diabetes by diet and lifestyle modification. *Journal of the American College of Nutrition* 2005; **24**: 310–9.
- Van Horn L, Obarzanek E, Friedman LA, Gernhofer N, Barton B. Children's adaptations to a fat-reduced diet: the Dietary Intervention Study in Children (DISC). *Pediatrics* 2005; **115**: 1723–33.
- Gardner CD, Coulston A, Chatterjee L, Rigby A, Spiller G, Farquhar JW. The effect of a plant-based diet on plasma lipids in hypercholesterolemic adults: a randomized trial. *Annals of Internal Medicine* 2005; **142**: 725–33.
- Fawzi WW, Msamanga GI, Spiegelman D, Wei R, Kapiga S, Villamor E, *et al.* A randomized trial of multivitamin supplements and HIV disease progression and mortality. *New England Journal of Medicine* 2004; **351**: 23–32.
- Kristal AR, Feng Z, Coates RJ, Oberman A, George V. Associations of race/ethnicity, education, and dietary intervention with the validity and reliability of a food frequency questionnaire: the Women's Health Trial Feasibility Study in Minority Populations. *American Journal of Epidemiology* 1997; **146**: 856–69.
- Willett W. *Nutritional Epidemiology*, 2nd edn. New York: Oxford University Press, 1998.
- Kristal AR, Beresford SA, Lazovich D. Assessing change in diet-intervention research. *American Journal of Clinical Nutrition* 1994; **59**(1 Suppl.): 185S–9S.
- Martinez ME, Marshall JR, Graver E, Whitacre RC, Woolf K, Rittenbaugh C, *et al.* Reliability and validity of a self-administered food frequency questionnaire in a chemoprevention trial of adenoma recurrence. *Cancer Epidemiology, Biomarkers & Prevention* 1999; **8**: 941–6.
- Simon MS, Lababidi S, Djuric Z, Uhley V, Depper J, Kresge C, *et al.* Comparison of dietary assessment methods in a low-fat dietary intervention program. *Nutrition and Cancer* 2001; **40**: 108–17.
- Thomson CA, Giuliano A, Rock CL, Ritenbaugh CK, Flatt SW, Faerber S, *et al.* Measuring dietary change in a diet intervention trial: comparing food frequency questionnaire and dietary recalls. *American Journal of Epidemiology* 2003; **157**: 754–62.
- Patterson RE, Kristal A, Rodabough R, Caan B, Lillington L, Mossavar-Rahmani Y, *et al.* Changes in food sources of dietary fat in response to an intensive low-fat dietary intervention: early results from the Women's Health Initiative. *Journal of the American Dietetic Association* 2003; **103**: 454–60.
- Sabaté J, Cordero-MacIntyre Z, Siapco G, Torabian S, Haddad E. Does regular walnut consumption lead to weight gain? *British Journal of Nutrition* 2005; **94**: 859–64.
- Knutsen SF, Fraser GE, Beeson WL, Lindsted KD, Shavlik DJ. Comparison of adipose tissue fatty acids with dietary fatty acids as measured by 24-hour recall and food frequency questionnaire in Black and White Adventists: the Adventist Health Study. *Annals of Epidemiology* 2003; **13**: 119–27.
- Knutsen SF, Fraser GE, Linsted KD, Beeson WL, Shavlik DJ. Comparing biological measurements of vitamin C, folate, alpha-tocopherol and carotene with 24-hour dietary recall information in nonhispanic blacks and whites. *Annals of Epidemiology* 2001; **11**: 406–16.
- Nutrition Coordinating Center, University of Minnesota. *Nutrition Data System for Research*. Minneapolis, MN: The Nutrition Coordinating Center, Division of Epidemiology, School of Public Health, University of Minnesota, 2000, 2003.
- Singh PN, Fraser GE, Knutsen SF, Lindsted KD, Bennett HW. Validity of a physical activity questionnaire among African-American Seventh-day Adventists. *Medicine and Science in Sports and Exercise* 2001; **33**: 468–75.
- Efron B, Tibshirani RJ. *An Introduction to the Bootstrap*. New York: Chapman and Hall, 1993.
- Almario RU, Vonghavaravat V, Wong R, Kasim-Karakas SE. Effects of walnut consumption on plasma fatty acids and lipoproteins in combined hyperlipidemia. *American Journal of Clinical Nutrition* 2001; **74**: 72–9.
- Chisholm A, Mann J, Skeaff M, Frampton C, Sutherland W, Duncan A, *et al.* A diet rich in walnuts favourably influences plasma fatty acid profile in moderately hyperlipidaemic subjects. *European Journal of Clinical Nutrition* 1998; **52**: 12–6.
- Potischman N, Carroll RJ, Iturria SJ, Mittl B, Curtin J, Thompson FE, *et al.* Comparison of the 60- and 100-item NCI-block questionnaires with validation data. *Nutrition and Cancer* 1999; **34**: 70–5.
- Subar AF, Thompson FE, Kipnis V, Midthune D, Hurwitz P, McNutt S, *et al.* Comparative validation of the Block, Willett, and National Cancer Institute food frequency questionnaires: the Eating at America's Table Study. *American Journal of Epidemiology* 2001; **154**: 1089–99.
- Ocke MC, Bueno-de-Mesquita HB, Pols MA, Smit HA, van Staveren WA, Kromhout D. The Dutch EPIC food frequency questionnaire. II. Relative validity and reproducibility for nutrients. *International Journal of Epidemiology* 1997; **26**(Suppl. 1): S49–58.
- Klipstein-Grobusch K, den Breeijen JH, Goldbohm RA, Geleijnse JM, Hofman A, Grobbee DE, *et al.* Dietary assessment in the elderly: validation of a semiquantitative food frequency questionnaire. *European Journal of Clinical Nutrition* 1998; **52**: 588–96.
- Yanek LR, Moy TF, Becker DM. Comparison of food frequency and dietary recall methods in African-American

- women. *Journal of the American Dietetic Association* 2001; **101**: 1361–4.
- 29 Flegal KM, Larkin FA. Partitioning macronutrient intake estimates from a food frequency questionnaire. *American Journal of Epidemiology* 1990; **131**: 1046–58.
- 30 Caan B, Ballard-Barbash R, Slattery ML, Pinsky JL, Iber FL, Mateski DJ, *et al.* Low energy reporting may increase in intervention participants enrolled in dietary intervention trials. *Journal of the American Dietetic Association* 2004; **104**: 357–66.
- 31 Siapco GS, Burns-Whitmore B, Haddad E, Cordero-MacIntyre Z. Weight change as a basis of determining energy intake reporting quality. *FASEB Journal* 2004; **18**: A860.
- 32 Block G. Invited commentary: another perspective on food frequency questionnaires. *American Journal of Epidemiology* 2001; **154**: 1103–4.